

# Processing of Analogy in the Thalamocortical Circuit

Yoonsuck Choe

Department of Computer Science, Texas A&M University  
3112 TAMU, College Station, TX 77843-3112  
Email: choe@tamu.edu

**Abstract**—The corticothalamic feedback and the thalamic reticular nucleus have gained much attention lately because of their integrative and modulatory functions. A previous study by the author suggested that this circuitry can process analogies (i.e., the *analogy hypothesis*). In this paper, the proposed model was implemented as a network of leaky integrate-and-fire neurons to test the *analogy hypothesis*. The previous proposal required specific delay and temporal dynamics, and the implemented network tuned accordingly functioned as predicted. Furthermore, these specific conditions turn out to be consistent with experimental data, suggesting that a further investigation of the thalamocortical circuit within the *analogical framework* may be worthwhile.

## I. INTRODUCTION

Understanding how cortical maps in the brain *interact* with each other to generate complex behavior is an important unsolved problem. Although we now know a lot more about the anatomical connectivity and physiology of the cortical maps than decades ago [2, 11], we still lack the understanding of how these maps work as an integrated system.

One key insight, from recent advances in neuroscience, can be gained from the thalamus. The thalamus is a centrally located nucleus in the brain with a high degree of feedforward and feedback connections to and from the cortex (see [23] for a review). The thalamus was previously thought of as a passive relay station for sensory-motor signals, however, this explanation was not satisfactory because of the existence of massive feedback from the cortex and a thin inhibitory network covering the thalamus called the thalamic reticular nucleus (TRN; see [6] for a review).

Previous work by the author suggested that the interplay between the thalamus, TRN, and the cortex may be implementing a function of analogy (the *analogy hypothesis*; [3, 4]). Analogy plays an important role in human perception and cognition [10, 12, 13, 16, 20], and its ability to cross domain boundaries may be critical in the integrative operation of cortical maps. Experimental observations by Crabtree and Isaac [6] nicely demonstrate the cross-modal nature of interaction in the thalamus and in the TRN, and these results can provide experimental grounds for the analogy hypothesis.

This paper presents the first computational implementation of the hypothetical model by the author initially described in [3, 4] and compare the results with the previous predictions. The following sections will briefly summarize the analogy hypothesis, then provide details of the implemented model, together with the results. The paper will conclude with a discussion about issues to be resolved, relevant related work, and future directions.




Fig. 1. **Analogy Through Active Completion and Filtering.** A simple brain with 6 detector neurons is shown. Environmental input comes in from the bottom (*World of Fruits*), and the neurons are linked with directed arcs that show the learned association between the detectors. From (a) to (c) shows the activation sequence of the brain in response to an analogy question “apple:orange::word-red:?” (a) Initial activation. (b) Active completion across relational links. (c) Remaining activity after filtering out input-driven cortical activity.

## II. ANALOGY THROUGH ACTIVE COMPLETION AND FILTERING

Neurons can be seen as processing information, i.e., producing output given a certain input. However, in this perspective, the output generated by neurons is *passive*, in the sense that they need further *interpretation* as all information or data does. A slight change of perspective allows us to view neurons as *active elements*, in the sense that neurons *actively invoke* other neurons. From this, it was shown that we can derive the function of analogy [3, 4]. The previous papers showed that *completion* and *filtering* are necessary for such active elements to implement the function of analogy. In the following, an example will briefly illustrate the proposed mechanism.

Let us suppose we have a simple brain with active neurons (or population of neurons) responding to specific inputs in the environment (Fig. 1). The simple brain has detectors for different input features including fruit objects (apple and orange), colors (red and orange color), and spoken words (word-red and word-orange). The neurons are *active*, i.e., they *invoke*

other neurons when they fire, and the invoking is done through relational (or associative) links which are learned through experience and embody frequently co-occurring events. For example, the apple-object detector has a strong connection to the color-red detector, etc.

Now consider the proportional analogy question `apple:orange::word-red:?`. Fig. 1 shows an activation sequence of the simple brain to the input question. Initially, the first 5 neurons are activated in response to the input (Fig. 1a). Next, through the relational links, other cortical neurons are invoked (*completion*; Fig. 1b). Last, after *filtering* for the purely cortex-driven activity in the cortex, only the word-orange detector remains active (Fig. 1c). We can see that word-orange is precisely the answer to the analogy question we posed in the beginning, and completion and filtering produced that answer.

However, simple filtering for only purely cortex-driven activity is insufficient for analogy answers containing items already present in the question. For example, consider the analogy question `big apple:small apple::big orange:?`. The answer is `small orange`, but both `small` and `orange` appeared in the question, thus the simple filtering as above will not work. In this case, relaxing the filtering criteria allows us to get to the answer: find *relatively less input-driven* cortical activities [3, 4].

In summary, simple analogies can be processed by active completion and filtering. In the next section, we will see how such completion and filtering can be neurally implemented.

### III. ACTIVE COMPLETION AND FILTERING IN THE THALAMOCORTICAL LOOP

How can active completion and filtering be implemented in the neural circuits of the brain? Previous work by the author proposed that the thalamus, thalamic reticular nucleus (TRN), and the cortex may be involved in completion and filtering [3, 4]. Fig. 2 shows a schematic diagram of the thalamocortical circuit based on known anatomy and physiology [6, 21]. In the following, we shall review how this circuit can carry out completion and filtering.

Cortico-cortical connections linking different maps in the cortex are ideally suited for active completion. However, a more difficult issue is how can filtering be done, i.e., how can input-driven cortical activity be distinguished from cortex-driven cortical activity? As suggested in [3, 4], the TRN is a promising location where such a filtering can occur.

The basic idea is that the reticular neurons receive both ascending thalamic input and descending cortical feedback, and reticular inhibition cancels out cortical feedback to the thalamic relays when both ascending and descending spikes are received at the TRN. On the other hand, when only descending (i.e., corticothalamic) feedback is received, TRN's inhibition on the thalamic relay is weak, and the relay neuron is allowed to fire in response to the cortical feedback, thus invoking the cortical neuron for the second time around.

Fig. 3 shows a detailed activation sequence of this mechanism. The example shows a case where only one of the




Fig. 2. **The Thalamocortical Circuit.** A schematic diagram of the thalamocortical circuit is shown (see [21] for a review). Solid edges with open arrows are excitatory connections, and dashed edges with closed arrows are inhibitory connections. Two thalamocortical loops ( $T_1$ - $R_1$ - $C_1$  and  $T_2$ - $R_2$ - $C_2$ ) are shown. The thalamus is a centrally located nucleus in the brain, and the dorsal part of it is covered with an inhibitory network of neurons called the Thalamic Reticular Nucleus (TRN). Ascending and descending connections all branch out and stimulate reticular neurons, and the reticular neurons send inhibitory connections to the thalamic relay neurons.

thalamocortical loops (loop1) received sensory input, thus demonstrating how *purely cortex-driven* activity can be singled out: The cortical feedback to the thalamus that survives the filtering is allowed to reactivate the cortex.

As for promoting the *relatively less input-driven* activity, we can think of a case when loop1 receives a strong sensory input and loop2 receives a weak sensory input. This time, both reticular neurons  $R_1$  and  $R_2$  will be highly activated, but due to the *disinhibition*<sup>1</sup> between the two,  $R_2$ , which is strongly inhibited by  $R_1$ , cannot cancel out cortical feedback from  $C_2$  to  $T_2$ . On the other hand,  $R_1$ , with its strong activity, will cancel out feedback from  $C_1$  to  $T_1$ .

#### A. Functional Requirements of the Proposed Circuit


There are specific assumptions that need to hold for the scenario described above to work:

- 1) reticular neurons must have a slow dynamics (Fig. 3b-d);
- 2) synaptic strength from the TRN to the thalamic relays must be weaker than those from the cortex to the thalamic relays (Fig. 3e);
- 3) either the cortico-cortical connections must be very fast or the corticothalamic feedback connections must be slow (or both), compared to each other (Fig. 3c-d); and
- 4) interaction between reticular neurons must be fast (Fig. 3d).

These conditions must hold due to the following functional requirements (in the same order as above):

- 1) reticular neurons need to retain the ascending excitation level to strongly inhibit the thalamic relay later when the cortical feedback comes around ( $R_1$  in Fig. 3b-d);

<sup>1</sup>Inhibition of an inhibitory neuron results in net excitation at the target of that neuron.



**Fig. 3. Analogical Completion and Filtering in the Thalamocortical Circuit.** An activation sequence of two thalamocortical loops (loop1 and loop2) are shown. Active parts of the circuit at each step are highlighted. (a) Initially, only  $T_1$  receives an afferent sensory input. (b)  $T_1$  invokes  $R_1$  and  $C_1$ . (c) The cortical neuron  $C_1$ , through fast connections, invokes another cortical neuron  $C_2$ .  $C_1$  also sends out feedback to  $R_1$  and  $T_1$ , but these connections are slow and the spike can only travel a short distance in the same time. Note that  $R_1$  retains the level of excitation because of its slow dynamics. (d) Cortical feedback from both  $C_1$  and  $C_2$  arrives at the TRN, and adds to the existing activity at TRN. Reticular neurons  $R_1$  and  $R_2$  inhibit each other through fast connections. (e) At the time the cortical feedback arrives at  $T_1$  and  $T_2$ , the reticular neurons exert inhibition on the thalamic relays.  $R_1$ , driven by both afferent input and cortical feedback exerts strong inhibition on  $T_1$ , effectively canceling out the cortical feedback from  $C_1$ . On the other hand,  $R_2$  was only driven by the cortical feedback, and it is not enough to cancel out feedback from  $C_2$  at the thalamic relay  $T_2$ . Thus  $T_2$  is permitted to fire again. As a result,  $C_2$  will be the only active neuron in the cortex in the next iteration.

- 2) reticular activity driven just by cortical feedback must not cancel out the feedback excitation from the cortex to the thalamic relay ( $R_2$  in Fig. 3e);
- 3) both input-driven cortical activity ( $C_1$ ) and cortical activity ( $C_2$ ) driven by that input-driven cortical activity must send feedback to the thalamus and TRN at approximately the same time (Fig. 3c–d); and
- 4) reticular neurons must rapidly adjust their activity level before inhibiting the thalamic relays ( $R_1$  and  $R_2$  in Fig. 3d).

Most of these conditions were described in [3, 4], but some of those that were only implicit in the earlier description are made more explicit here.

As it turns out, all of these conditions have experimental support (in the same order as above):

- 1) reticular neurons activate and deactivate on a slow timescale compared to thalamic relays [5, 14];
- 2) reticular neurons are harder to depolarize than thalamic relays [14]<sup>2</sup>;
- 3) corticothalamic feedback connections are unmyelinated (i.e., very slow) [25]; and

<sup>2</sup>Sherman and Guillery make a cautionary note against a similar interpretation of the phenomenon regarding low-threshold  $Ca^{2+}$  spikes in the reticular neurons [23], pp.138–139.

- 4) gap junctions have been found between reticular neurons [18], suggesting that the interaction between reticular neurons may be rapid.<sup>3</sup>

In the following, the conditions listed above will be tested in a computational implementation of the model.

#### IV. MODEL DESCRIPTION

A network of leaky integrate-and-fire neurons [9] was constructed to test the *analogy hypothesis* summarized in the previous section.

Six neurons of three types (T: thalamic relay; R: thalamic reticular neuron; C: cortical neuron) were connected according to the diagram in Fig. 2 (with the additional connection from  $C_2$  to  $C_1$ ). For each neuron  $i$ , the membrane potential  $V_i$  evolved according to the following dynamic equation:

$$C_i \frac{dV_i}{dt} = I_i(t) - \frac{V_i}{R_i}, \quad (1)$$

where  $C_i$  is the membrane capacitance,  $R_i$  the resistance, and  $I_i(t)$  the input contribution to neuron  $i$  at time  $t$ . When  $V_i$  reaches a threshold value  $\theta_i$ , a spike is generated and  $V_i$  is reset to 0.0. A spike generated by a presynaptic neuron  $j$  results in a postsynaptic potential (PSP)  $s_{ij}$  at a target neuron  $i$ , which is set to 1.0 at the moment the spike is received and is decayed over time as follows:

$$\frac{ds_{ij}}{dt} = -\frac{s_{ij}}{\tau_i}, \quad (2)$$

where  $\tau_i$  is the time constant of the PSP in the neuron  $i$ .

The input contribution  $I_i(t)$  to the neuron  $i$  at time  $t$  is defined as follows:

$$I_i(t) = \sum_{j \in \mathcal{N}_i} w_{ij} s_{ij}(t - \delta_{ij}), \quad (3)$$

where  $\mathcal{N}_i$  is the set of neurons sending spikes to neuron  $i$  (see Fig. 2);  $w_{ij}$  is the connection weight from neuron  $j$  to  $i$  (the sign is negative if  $j$  is an inhibitory neuron); and  $s_{ij}(t - \delta_{ij})$  is the PSP generated in neuron  $i$  by a spike from neuron  $j$  with a conduction delay of  $\delta_{ij}$ . See Section V, Tables I and II for the exact parameter values.

#### V. EXPERIMENTS AND RESULTS

Three experiments were conducted with the model described above to test the assumptions and predictions in the hypothetical model (Fig. 2; [3, 4]). The experiments tested if the model tuned according to the conditions listed in Section III-A can filter out input-driven or less input-driven cortical activity and just leave either the purely cortex-driven or relatively more cortex-driven activity in the cortex.

##### A. General Experimental Setup

Tables I and II below list the neuron and connection parameters (the units are arbitrary). The parameters are fairly uniform, except for the figures in **bold-type** indicating a deviation from the default parameters.

<sup>3</sup>It is controversial whether gap junctions can carry out disinhibition as required in here. However, the existence of gap junctions shows that reticular neurons need to communicate at a high speed, suggesting that other connections between reticular neurons may have to be fast as well.




Fig. 4. **Thalamus- vs. Cortex-Driven Activity.** Membrane potential traces for a thalamic relay (T), a reticular neuron (R), and a cortical neuron (C) in a *single* loop are shown from bottom to top in each panel. The x-axis is time and the y-axis is the membrane potential. A depolarizing current of magnitude 1.0 and duration 1 was either injected in (a) the thalamic relay, emulating an input-driven condition, or in (b) the cortical neuron, emulating a cortex-driven condition. Only for the cortex-driven case (b), the initial cortical burst of activity can reactivate the cortical neuron through the corticothalamic loop. Note that the plot shows the  $V_i$  trace (equation 1) with spikes added at the moment of threshold crossing. Note that (a) and (b) are from two independent experiments.

TABLE I  
NEURON PARAMETERS

Parameter	Thal. Relay ( $T_i$ )	TRN ( $R_i$ )	Cortex ( $C_i$ )
Capacitance $C_i$	0.3	<b>0.6</b>	0.3
Resistance $R_i$	3.0	3.0	3.0
Threshold $\theta_i$	0.25	0.25	0.25
PSP time constant $\tau_i$	0.05	0.05	0.05

TABLE II  
CONNECTION PARAMETERS

Weight $w_{ij}$	$T_i$	$R_i$	$C_i$	Delay $\delta_{ij}$	$T_i$	$R_i$	$C_i$
$T_j$		1.0	1.0	$T_j$		2.0	2.0
$R_j$	<b>0.5</b>	<b>10.0</b>		$R_j$	2.0	<b>0.2</b>	
$C_j$	1.0	1.0	1.0	$C_j$	<b>4.0</b>	2.0	<b>0.2</b>

The deviations are not arbitrary, and are specifically required as discussed in Section III-A (in the same order):

- 1) the membrane capacitance  $C_i$  of a reticular neuron  $R_i$  must be large ( $= 0.6$ , twice the default value) so that the membrane has slow dynamics;
- 2) the connection weight  $w_{ij}$  from a reticular neuron  $R_i$  to a thalamic relay  $T_i$  must be small ( $= 0.5$ , half the default value);
- 3) the conduction delay  $\delta_{ij}$  from a cortical neuron  $C_j$  to a thalamic relay  $T_i$  must be large ( $= 4.0$ , twice the default value); the conduction delay  $\delta_{ij}$  from a cortical neuron  $C_j$  to another cortical neuron  $C_i$  must be very small ( $= 0.2$ , 1/10th the default value); and
- 4) the conduction delay  $\delta_{ij}$  from a reticular neuron  $R_j$  to  $R_i$  must be small ( $= 0.2$ , 1/10th the default value).

One exception is the connection weight  $w_{ij}$  from  $R_j$  to  $R_i$ , the implication of which has to be investigated further.

The same set of parameters were used without modification in all the experiments to follow.

### B. Experiment 1: Input-Driven vs. Cortex-Driven Activity in a Single Loop

To independently test the function of a *single* thalamocortical loop, depolarizing currents were injected at two different

locations of the loop in two separate experiments: (1) the thalamic relay T and (2) the cortical neuron C. The thalamic injection emulated the input-driven condition, and the cortical injection the cortex-driven condition. The results are shown in Fig. 4.

In the input-driven case (Fig. 4a), the cortex activates at time  $t = 3$  in response to the thalamic drive, but the corticothalamic feedback at time  $t = 7$  is canceled out (i.e., filtered out) by the reticular inhibition. As a result, the cortical neuron fails to reactivate.

In contrast, for the cortex-driven case (Fig. 4b), the corticothalamic feedback to T at time  $t = 5$  is strong enough to survive the weak reticular inhibition and successfully reactivates the cortex at time  $t = 7$ .

### C. Experiment 2: Input vs. No-Input Condition in a Pair of Loops

In this experiment, a pair of thalamocortical loops was simulated to test whether *purely cortex-driven* activity can be singled out in the model. The two loops were connected as in Fig 2 with an addition of reciprocal cortico-cortical connection from  $C_2$  to  $C_1$ . Input current was only injected to loop1 thalamic relay  $T_1$ . The results are shown in Fig. 5.

For the initially input-driven loop (loop1; Fig. 5a), the final cortical burst of activity at time  $t = 9$  is unable to reactivate the cortex. However, for the initially cortex-driven loop (loop2; Fig. 5b), the cortical burst at time  $t = 9$  is able to reactivate the cortex at  $t = 16$  through the corticothalamic feedback. Thus, the model demonstrates selectivity for purely cortex-driven activity.<sup>4</sup>

### D. Experiment 3: Strong- vs. Weak-Input Condition in a Pair of Loops

The setup in this experiment was the identical to Section V-C, except for the input condition. For this experiment, loop1 thalamic relay  $T_1$  was injected with a depolarizing current of magnitude 2.0, and  $T_2$  of loop2 was injected with a current, but

<sup>4</sup>Note that  $C_1$  reactivates once but not twice. The reason it reactivates is because of the recursive cortico-cortical activation coming from  $C_2$ .




Fig. 5. **Input vs. No-Input Condition.** Membrane potential traces for neurons in two connected thalamocortical loops are shown, one in (a) and the other in (b). The two loops are wired as shown in Fig. 2, with the addition of a reciprocal cortico-cortical connection from  $C_2$  to  $C_1$ . The voltage traces for the neurons in the loops are numbered accordingly. (a) A depolarizing current of magnitude 1.0, duration 1 was injected in  $T_1$ . (b) No current was injected anywhere in the loop, thus all activities were initially driven by the cortico-cortical connection from  $C_1$  to  $C_2$  at time  $t = 3$ . In the end, only the cortex-driven cortical activity ( $C_2$ ) is able to reactivate the cortex through feedback to the thalamus (panel b). Note that the above are results from a single experiment.




Fig. 6. **Strong- vs. Weak-Input Condition.** Membrane potential traces for two thalamocortical loops with an identical setup as in Fig. 5 is shown. The only difference was the magnitude of depolarizing current injected in loops 1 and 2. (a) A depolarizing current of magnitude 2.0, duration 1 was injected in  $T_1$ . (b) A depolarizing current of magnitude 1.0, duration 1 was injected in  $T_2$ . Thus, both loops were input-driven, but to a different degree. In the end, only the less input-driven cortical activity ( $C_2$ ) is able to reactivate the cortex through feedback to the thalamus.

with a lower magnitude of 1.0. Thus, a strongly input-driven vs. weakly input-driven condition was setup to test whether *relatively less input-driven* cortical activity can be promoted in the model. The results are shown in Fig. 6.

The results are similar to those in Section V-C. Cortical activity in the more input-driven loop1 is unable to reactivate the cortex twice (Fig. 6a), but the less input-driven loop2 is able to reactivate the cortex twice ( $t = 16$ ; Fig 6b). Again, these results show that the model has selectivity for less input-driven cortical activity.

Note that in this case, disinhibition between reticular neurons play an important role in allowing loop2 to reactivate the cortex, despite the fact that loop2 was also input-driven: Loop1 reticular neuron  $R_1$  fire two times more compared to  $R_2$  of loop2, and when they fire, they inhibit  $R_2$  ( $t = 3, 5,$  and  $12$ ), thus weakening the inhibition exerted by  $R_2$ .

*E. Summary*

In summary, the thalamocortical model functioned as predicted by the *analogy hypothesis*, with a fixed set of parameters derived from physiological considerations. The model was successful in detecting and promoting (1) purely cortex-driven

cortical activity, and (2) relatively less input-driven cortical activity, which are requirements for the processing of analogy.

VI. DISCUSSION

Two major emphases of this paper are *activeness* and the integrative role of *analogy* in brain function. The two rather abstract notions turn out to have a firm biological ground (i.e., the thalamocortical loop), and through this connection we can begin to take a fresh look at how the diverse cortical maps can interact to give rise to an integrated behavior. Action [1, 15], analogy [10, 12, 13, 16, 20], and the attentive and integrative role of the thalamocortical circuit [6, 17, 19, 21–23] are all related in that respect, and a collective effort in understanding the relationship between these areas will become necessary.

The current model can only address a limited range of analogy problems [3, 4], and these issues need to be resolved. The most prominent issues are that of structured analogy and that of temporal or spatial order. Investigations into these issues will inevitably involve the prefrontal cortex [8], which is believed to be dealing with sequences of events.

Another issue not addressed in the previous papers [3, 4] is that of synchrony [24, 26]. Thalamus plays an important

role in synchronization [19], and how analogical processes can interact with synchronized populations of neurons will become an important issue.

## VII. CONCLUSION

The current work computationally tested an earlier proposal by the author that the thalamocortical circuit may be performing analogies across cortical maps. A network of integrate-and-fire neurons was built and tuned based on functional and physiological considerations. The results showed that *active completion* and *filtering* for less input-driven activity, which forms the basis of analogy, arise in the model. These results suggest that further investigation into the thalamocortical mechanism of analogy may be worthwhile.

## ACKNOWLEDGMENTS

I would like to thank Bard Ermentrout for his help with XPPAUT [7], in which the model was implemented. This research was supported in part by Texas A&M University, by the Texas Higher Education Coordinating Board ARP/ATP program (#000512-0217-2001), and by the National Institute of Mental Health Human Brain Project (#1R01-MH66991).

## REFERENCES

- [1] Arbib, M. A. (2003). Schema theory. In Arbib, M. A., editor, *The Handbook of Brain Theory and Neural Networks*, 993–998. Cambridge, MA: MIT Press. Second edition.
- [2] Bosking, W. H., Zhang, Y., Schofield, B., and Fitzpatrick, D. (1997). Orientation selectivity and the arrangement of horizontal connections in tree shrew striate cortex. *Journal of Neuroscience*, 17(6):2112–2127.
- [3] Choe, Y. (2002). Second order isomorphism: A reinterpretation and its implications in brain and cognitive sciences. In Gray, W. D., and Schunn, C. D., editors, *Proceedings of the 24th Annual Conference of the Cognitive Science Society*, 190–195. Erlbaum.
- [4] Choe, Y. (2003). Analogical cascade: A theory on the role of the thalamo-cortical loop in brain function. *Neurocomputing*. In press.
- [5] Coulter, D. A., Huguenard, J. R., and Prince, D. A. (1989). Calcium currents in rat thalamocortical relay neurons: Kinetic properties of the transient, low-threshold current. *Journal of Physiology*, 414:587–604.
- [6] Crabtree, J. W., and Isaac, J. T. (2002). Intrathalamic pathways allowing modality-related and cross-modality switching in the dorsal thalamus. *The Journal of Neuroscience*, 22:8754–8761.
- [7] Ermentrout, B. (2002). *Simulating, Analyzing, and Animating Dynamical Systems: A Guide to XPPAUT for Researchers and Students*. Philadelphia, PA: SIAM.
- [8] Fuster, J. M. (1997). *The Prefrontal Cortex: Anatomy, Physiology, and the Frontal Lobe*. Lippencott-Raven. Third edition.
- [9] Gabbini, F., and Koch, C. (1999). Principles of spike train analysis. In Koch, C., and Segev, I., editors, *Methods in Neural Modeling*, chapter 7, 313–360. MIT Press.
- [10] Gentner, D., Holyoak, K. J., and Kokinov, B. N., editors (2001). *The Analogical Mind: Perspectives from Cognitive Science*. Cambridge, MA: MIT Press.
- [11] Gilbert, C. D., and Wiesel, T. N. (1989). Columnar specificity of intrinsic horizontal and corticocortical connections in cat visual cortex. *Journal of Neuroscience*, 9:2432–2442.
- [12] Hofstadter, D. R. (1995). *Fluid Concepts and Creative Analogies: Computer Models of the Fundamental Mechanisms of Thought*. Basic Books.
- [13] Holland, J. H., Holyoak, K. J., Nisbett, R. E., and Thagard, P. R. (1986). *Induction: Process of Inference, Learning, and Discovery*. Cambridge, MA: MIT Press.
- [14] Huguenard, J. R., and McCormick, D. A. (1992). A novel t-type current underlies prolonged  $Ca^{2+}$ -dependent burst firing in GABAergic neurons of rat thalamic reticular nucleus. *Journal of Neuroscience*, 12:3804–3817.
- [15] Hurley, S. (2001). Perception and action: Alternative views. *Synthese*, 129:3–40.
- [16] Kanerva, P. (1998). Dual role of analogy in the design of a cognitive computer. In Holyoak, K., Gentner, D., and Kokinov, B., editors, *Advances in Analogy Research: Integration of Theory and Data from the Cognitive, Computational, and Neural Sciences*, 164–170.
- [17] LaBerge, D. (1995). *Attentional Processing*. Cambridge, MA: Harvard University Press.
- [18] Landisman, C. E., Long, M. A., Beierlein, M., Deans, M. R., Paul, D. L., and Connors, B. W. (2002). Electrical synapses in the thalamic reticular nucleus. *Journal of Neuroscience*, 22:1002–1009.
- [19] Linás, R. R., and Ribary, U. (1994). Perception as an oneiric-like state modulated by the senses. In Koch, C., and Davis, J. L., editors, *Large Scale Neuronal Theories of the Brain*, chapter 6, 111–124. Cambridge, MA: MIT Press.
- [20] Mitchell, M. (1993). *Analogy-Making as Perception: A Computer Model*. Cambridge, MA: MIT Press.
- [21] Mumford, D. (1995). Thalamus. In Arbib, M. A., editor, *The Handbook of Brain Theory and Neural Networks*, 153–157. Cambridge, MA: MIT Press.
- [22] Newman, J., Baars, B. J., and Cho, S.-B. (1997). A neural global workspace model for conscious attention. *Neural Networks*, 10:1195–1206.
- [23] Sherman, S. M., and Guillery, R. W. (2001). *Exploring the Thalamus*. San Diego, CA: Academic Press.
- [24] Singer, W., and Gray, C. M. (1995). Visual feature integration and the temporal correlation hypothesis. *Annual Review of Neuroscience*, 18:555–586.
- [25] Tsumoto, T., Creutzfeldt, O. D., and Legendy, C. R. (1978). Functional organization of the corticofugal system from visual cortex to lateral geniculate nucleus in the cat. *Experimental Brain Research*, 32:345–364.
- [26] Wang, D. L. (1995). Emergent synchrony in locally coupled neural oscillators. *IEEE Transactions on Neural Networks*, 6:941–948.